

Relic HII Regions and Radiative Feedback at High-Redshifts

Andrei Mesinger^{1*}, Greg L. Bryan², & Zoltán Haiman²

¹*Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA*

²*Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027, USA*

13 September 2009

ABSTRACT

UV radiation from early astrophysical sources could have a large impact on subsequent star formation in nearby protogalaxies, and in general on the progress of cosmological reionization. Theoretical arguments based on the absence of metals in the early Universe suggest that the first stars were likely massive, bright, yet short-lived, with lifetimes of a few million years. Here we study the radiative feedback arising from such stars using hydrodynamical simulations with transient UV backgrounds (UVBs) and persistent Lyman-Werner backgrounds (LWBs) of varying intensity. We extend our prior work in Mesinger et al. (2006), by studying a more typical region whose proto-galaxies form at lower redshifts, $z \sim 13$ –20, in the epoch likely preceding the bulk of reionization. We confirm our previous results that feedback in the relic HII regions resulting from such transient radiation, is itself transient. Feedback effects dwindle away after $\sim 30\%$ of the Hubble time, and the same critical specific intensity of $J_{\text{UV}} \sim 0.1 \times 10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ separates positive and negative feedback regimes. This suggests that overall feedback is fairly insensitive to the large-scale environment, overdensity, and redshift-dependent halo parameters, and can accurately be modeled in this regime with just the intensity of the impinging UVB. Additionally, we discover a second episode of eventual positive feedback in halos which have not yet collapsed when their progenitor regions were exposed to the transient UVB. When exposed to the transient UVB, this gas suffers relatively little density depletion but a significant enhancement of the molecular hydrogen abundance, thus resulting in net positive feedback. This eventual positive feedback appears in all runs, regardless of the strength of the UVB. However, this feedback regime is very sensitive to the presence of Lyman-Werner radiation, and notable effects disappear under fairly modest background intensities of $J_{\text{LW}} \gtrsim 10^{-3} \times 10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$, assuming the region is optically thin for LW photons. Nevertheless, when exposed to the same LWB, halos inside relic HII regions always have a higher H_2 abundance and shorter cooling times than halos outside relic HII regions, allowing gas to cool faster once it finally begins to collapse onto the halo. We conclude that UV radiative feedback in relic HII regions, although a complicated process, seems unlikely to have a major impact on the progress of cosmological reionization, provided that present estimates of the lifetime and luminosity of a Pop III star are accurate. More likely is that the build-up of the LWB ultimately governs the feedback strength until a persistent UV background can be established.

Key words: cosmology: theory – early Universe – galaxies: high-redshift – evolution

1 INTRODUCTION

Semi-analytic models and numerical simulations predict that the first astrophysical objects formed at redshifts $z \gtrsim 20$. They form out of metal-free gas with inefficient fragmentation, thus numerical simulations predict that these first objects could be very massive ($M_* \gtrsim 100 M_\odot$), so-called ‘Population III’ (Pop III) stars (Abel et al. 2002; Bromm et al. 2002; Yoshida et al. 2008). These stars would be short-lived, with lifetimes of ~ 3 Myr, but could produce ten times more ionizing photons per baryon than regular

Population II stars (Schaerer 2002, 2003). Also, such star formation could have been somewhat synchronized, given that these early objects formed in highly biased and clustered environments. Hence, Pop III stars could have a significant impact on subsequent generations of objects.

Indeed Pop III stars can strongly affect the progress of cosmological reionization (Haiman & Holder 2003; Cen 2003b,a; Wyithe & Loeb 2003), although their contribution is virtually unknown from first principles and is generally applied to simulations with a “tuning knob” approach. Thus ionizing photons from the first stars and lack thereof have been used to explain both early (e.g. Cen 2003a) and fairly late (e.g. Haiman & Bryan 2006) reion-

* Hubble Fellow; e-mail: mesinger@astro.princeton.edu

ization, depending on which scenario was favored when the works were published. Therefore, it is quite important to gain physical insight into how the radiation of the first stars impacted their surroundings.

Radiative feedback can be either positive or negative, in that it can enhance or suppress subsequent star-formation. Positive feedback can result when the enhanced free-electron fraction from ionizing photons (e.g. Oh & Haiman 2002) or hydrodynamical shocks (Shapiro & Kang 1987) catalyzes the formation of molecular hydrogen (H_2). If the ionizing background “turns-off” (resulting in so-called “relic” HII regions), H_2 cooling can provide the dominant cooling channel at high-densities and low temperatures. Conversely, negative feedback can result from heating by ionizing radiation which can photo-evaporate gas in low-mass halos (Efstathiou 1992; Barkana & Loeb 1999; Gnedin 2000; Shapiro et al. 2004; Dijkstra et al. 2004; Mesinger & Dijkstra 2008). Also, an active background of Lyman-Werner (LW) radiation (with photon energies in the 11.18–13.6eV range) can dissociate H_2 , thus decreasing the gas’s cooling capabilities (e.g., Haiman et al. 1997, 2000; Ciardi et al. 2000; Machacek et al. 2001; Wise & Abel 2007; O’Shea & Norman 2008).

Indeed, simulations find radiative feedback to be nuanced at very high redshifts. Positive feedback dominates in flash ionized gas (O’Shea et al. 2005), or gas exposed to a weak transient ultraviolet background (UVB) (Mesinger et al. 2006, hereafter MBH06) such as might be present close to the edges of HII regions (Ricotti et al. 2002; Kuhlen & Madau 2005). On the other hand, negative feedback in relic HII regions can occur in regions closer to the Pop III star (MBH06; Susa & Umemura 2006; Ahn & Shapiro 2007; Yoshida et al. 2007). Furthermore, radiative transfer effects can also impact the strength and sign of feedback (e.g. Iliiev et al. 2005; Susa & Umemura 2006; Abel et al. 2007; Whalen et al. 2008; Wise & Abel 2008b). Because of variations in halo mass, collapse redshift, central baryon density, stellar spectrum, and distance to the ionizing source, it is difficult to accurately deal with the large parameter space governing photoevaporation.

In MBH06, we attempted to tackle this issue with a statistical approach. By giving up on simulating “realistic” (as much as this is possible given present uncertainties) relic HII regions left behind by deceased Pop III stars, we were able to explore larger swaths of parameter space and have a fairly large sample of second-generation objects. We did this by applying various uniform ultraviolet (UV) and LW backgrounds on our entire simulation box, and statistically comparing the resulting evolution against the fiducial run without radiation. Although this approach clearly does not realistically treat the photoionization around a particular source, it might provide a good statistical description of an ensemble of reionization studies. Clearly, at the onset of early reionization not all halos lie within HII regions, as they do in our models. Our primary purpose is not to simulate early reionization or the photoevaporation of well-formed Pop III halos in detail (which would require radiative transfer) but to evaluate the impact of relic HII regions on the formation of halos at later times. Our uniformly illuminated boxes provide an ensemble of evaporated halos at high redshift that statistically sample the effects of early photoionization on the assembly of halos and formation of cold dense gas at lower redshifts.

However, since our analysis in MBH06 focused on a highly biased region which went non-linear at a high redshift, we could not follow the evolution to redshifts much lower than $z \sim 20$. This meant that we could only extrapolate the observed trends suggesting that the feedback was transient. Oh & Haiman (2003) claim that, even in the presence of a weak LWB, excess entropy will even-

tually suppress star formation in protogalaxies forming inside relic HII regions. In MBH06, we were unable to conclusively verify or refute this claim.

In this work, we extend the study by MBH06 by analyzing radiative feedback in a less-biased region, whose typical protogalaxies form at lower redshifts, $z \sim 13$ –20. Given that the 5-yr *WMAP* polarization data place a constraint on the reionization redshift (assuming instantaneous reionization) of $z = 11.0 \pm 1.4$ (Dunkley et al. 2008), such cosmological regions are likely to host the majority of feedback effects from Pop III stars, before a persistent UVB and/or a strong LWB become entrenched (e.g. Haiman et al. 1996). We apply the same statistical approach as in MBH06, and study various combinations of transient UVBs and persistent LWBs. As such, we attempt to provide a framework for analytically incorporating such radiative feedback into semi-analytic, and large-scale numerical and semi-numerical studies.

In §2 we enumerate and describe the simulations which are used in this work. In §3, we study the initial radiative feedback caused by a transient UVB, adding a persistent LWB in §3.1. Then in §4, we describe a related mechanism that results in eventual positive feedback, regardless of the strength of the UVB. In §5, we discuss the impact of neglecting self-shielding. Finally, we offer our conclusions in §6.

Throughout this paper, we adopt the background cosmological parameters $(\Omega_\Lambda, \Omega_M, \Omega_b, n, \sigma_8, H_0) = (0.7, 0.3, 0.047, 1, 0.92, 70 \text{ km s}^{-1} \text{ Mpc}^{-1})$, consistent with the measurements of the power spectrum of CMB temperature anisotropies by the first year of data from the *WMAP* satellite (Spergel et al. 2003). Although the 5-yr data prefers a slightly lower value of $\sigma_8 = 0.82$ (Dunkley et al. 2008; Komatsu et al. 2008) which somewhat delays structure formation, we keep the cosmology the same as in MBH06, to facilitate direct comparison. Unless stated otherwise, we quote all quantities in comoving units.

2 SIMULATIONS

We use the Eulerian adaptive mesh refinement (AMR) code Enzo, which is described in greater detail elsewhere (Bryan 1999; Norman & Bryan 1999). Our simulation volume is $1 (h^{-1} \text{ Mpc})^3$, initialized at $z_{\text{init}} = 99$ with density perturbations drawn from the Eisenstein & Hu (1999) power spectrum. Our root grid is 128^3 . We have two additional static levels of refinement inside a central $0.25 h^{-1} \text{ Mpc}$ cube. In addition, grid cells inside the central region are allowed to dynamically refine so that the Jeans length is resolved by at least 4 grid zones and no grid cell contains more than 4 times the initial gas mass element. Each additional grid level refines the mesh length of the parent grid cell by a factor of 2. We allow for a maximum of 10 levels of refinement inside the refined central region, granting us a spatial resolution of $7.63 h^{-1} \text{ pc}$. This comoving resolution translates to $0.36 h^{-1}$ proper pc at $z = 20$.

As stated above, our simulation runs were set up to facilitate comparison with MBH06. Users interested in details of the simulations are encouraged to consult MBH06 and Machacek et al. (2001). There is one notable difference between our runs here and those in MBH06: our central refined region for these runs is *not* centered on the highest density region of the box. Instead it is chosen to be more typical of the regions expected to host the bulk of halos which form prior to any significant cosmological reionization. Specifically, the density inside our central $(0.25 h^{-1} \text{ Mpc})^3$ corresponds to a 0.75σ mass fluctuation of an equivalent spherical volume (in MBH06, we studied a 2.4σ region). The less overdense

region allows us to extend the analysis of MBH06 to lower redshifts, and test the robustness of the conclusions of MBH06 on a less-biased and more typical region.

As shown in Table 1, we have performed four different runs without a LW background, distinguished by the duration or amplitude of the assumed UVB, and eight additional runs that include an additional constant LW background. Again, these runs are analogous to those in MBH06. For the UV radiation we assume an isotropic background flux with a $T = 2 \times 10^4$ K blackbody spectral shape, normalized at the hydrogen ionization frequency, $h\nu_H = 13.6$ eV. This spectrum is softer than the $\sim 8 \times 10^4$ K blackbody spectrum predicted to be produced by Pop III stars (e.g. Schaerer 2002), which leads us to find a lower reionization temperature (by about a factor of two) than we would have found with a harder spectrum. This results in outflow velocities (and therefore halo evacuation times) which are also too low by this factor.

The values of J_{21}^{UV} are shown in Table 1 in units of $10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$. The NoUVB run contains no UV radiation, and serves mainly as a reference run. The Heat0.08 and Heat0.8 runs include a UVB with $J_{21}^{UV} = 0.08, 0.8$, respectively. The value of $J_{21}^{UV} = 0.08$ was chosen to correspond to the mean UV flux expected inside a typical HII region surrounding a primordial star (e.g., Alvarez et al. 2006). As we do not include dynamically expanding HII regions in our code, the Heat0.8 and Flash (in which the UVB is turned on for a negligibly short duration) runs can be viewed as extremes, corresponding to conditions close to the center and close to the edge of the HII region, respectively. More generally, studying a range of values of J_{UV} is useful, since the radiative properties of high-redshift sources are uncertain. In the latter two runs, the UVB is turned on at $z_{UVB, \text{on}} = 25$ and turned off at $z_{UVB, \text{off}} = 24.64$. This redshift range corresponds to a typical theoretical stellar lifetime, ~ 3 Myr, of a $\sim 100 M_\odot$ primordial (Pop-III) star (Schaerer 2002). The flash ionization run, Flash, instantaneously sets the gas temperature to $T=15000$ K and the hydrogen neutral fraction to $x_{\text{HI}} = 10^{-3}$ throughout the simulation volume, but involves no heating thereafter. Again, our adopted temperature is probably too low by a factor of two compared to gas heated by Pop III stars. Finally, the eight runs in the bottom half of the Table repeat pairs of the NoUVB and Heat0.8 runs with four different constant LW backgrounds ($J_{21}^{\text{LW}} = 10^{-4}, 0.001, 0.01$, and 0.1 , normalized at 12.87eV in units of $10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$, and assumed to be frequency-independent within the narrow LW band).

We obtain the dark matter (DM) and baryonic mass of halos, in the same manner as MBH06. Similarly, we use the fraction of total gas within the virial radius which is cold and dense (CD), f_{cd} as a proxy for the gas reservoir available to form stars. By cold, we mean gas whose temperature is $< 0.5 T_{\text{vir}}$, where T_{vir} is the halo’s virial temperature (for how virial temperatures are associated with halos in the simulation, see Machacek et al. 2001). By dense, we mean gas whose density is $> 10^{19} M_\odot \text{ Mpc}^{-3} \approx 330 \text{ cm}^{-3}$, roughly corresponding to the density at which the baryons become important to the gravitational potential in the core, taken to be an immediate precursor to primordial star formation (Abel et al. 2002).

Additionally, we discount halos which have been substantially contaminated by the large (low-resolution) DM particles originally outside of our refined region which enter it as it becomes nonlinear. Specifically, we remove from our analysis halos with an average DM particle mass greater than $\sim 140\%$ of the refined region’s DM mass resolution, $747 M_\odot$. However, since there are small stochastic variations in the amount of contamination between the runs, we first

Table 1. Summary of Simulation Runs

Run Name	J_{21}^{UV}	$z_{UVB, \text{on}}$	$z_{UVB, \text{off}}$	J_{21}^{LW}
Runs without a LW background				
NoUVB	0	NA	NA	0
Flash	–	25	25	0
Heat0.08	0.08	25	24.62	0
Heat0.8	0.8	25	24.62	0
Runs with a LW background				
LW 10^{-4} _NoUVB	0	NA	NA	10^{-4}
LW 10^{-4} _Heat0.8	0.8	25	24.62	10^{-4}
LW 10^{-3} _NoUVB	0	NA	NA	10^{-3}
LW 10^{-3} _Heat0.8	0.8	25	24.62	10^{-3}
LW 10^{-2} _NoUVB	0	NA	NA	0.01
LW 10^{-2} _Heat0.8	0.8	25	24.62	0.01
LW0.1_NoUVB	0	NA	NA	0.1
LW0.1_Heat0.8	0.8	25	24.62	0.1

apply this cut-off in the NoUVB run, and then remove the same halos from the analysis of the other runs. This way, we study the same halos in each run.

Another possible source of contamination arises from closely separated halos. If some CD gas belonging to a halo is within another halo’s virial radius (most likely in the process of merging), the other halo could undeservedly be flagged as containing CD gas as well. To counteract this, we set $f_{\text{cd}} = 0$ for all halos without CD gas in their inner-most spherical shell. CD gas invariably first forms close to the halo’s core, where the cooling times are the shortest; thus halos which contain CD gas at large radii but none in their core are most likely in the process of merging with a halo with genuine CD gas. Note that this algorithm is different than the one used in MBH06, where we discriminated based on mass and separation between halos. In this work such contamination was more prevalent and so we devised what proved to be a more robust means of discrimination.

Finally, we note that we do not include HD cooling in these simulations. As is shown in McGreer & Bryan (2008) and Yoshida et al. (2007), HD cooling is important in primordial halos only in the density range $10^2 - 10^6 \text{ cm}^{-3}$, where H_2 can bring the temperature below about 200 K. Therefore it does not change the amount of CD gas, our primary diagnostic (although it would lead to relatively small changes in the predicted masses of the resulting stars).

3 TRANSIENT FEEDBACK FOLLOWING A TRANSIENT UVB

We first wish to confirm whether our conclusions in MBH06 also apply to this new sample of halos, which form at later times in a less overdense region of the universe. In MBH06, we found the following. The UVB ionizes and heats the gas in our simulation box.¹ Gas inside halos which used to be at or close to hydrostatic equilibrium feels an increased pressure gradient from the rise in temperature, resulting in an outward-moving pressure shock. Once the UVB is

¹ It is important to note that we are operating in the optically thin limit; the accuracy and impact of that approximation is discussed in section 5.




Figure 1. Spherically averaged radial profiles of the same halo in the NoUVB (solid lines), Heat0.08 (dashed lines) and Heat0.8 (dotted lines) simulation runs. This halo was first able to form CD gas at $z = 20, 21, 19$ in the NoUVB, Heat0.08, and Heat0.8 runs, respectively. The top and bottom pairs of figures show snapshots at $z = 24.62, 23, 21, 19$ (left to right). The halo mass in the NoUVB run increases from $M = 4.2 \times 10^5 M_\odot$ at $z = 24.62$, to $M = 1.4 \times 10^6 M_\odot$ at $z = 19$. All quantities are shown in proper (not comoving) units. The *Upper panels* show the hydrogen density, mass-weighted gas temperature, gas cooling time, and radial velocity (clockwise from upper left). The *Bottom panels* show mass fractions of HI, HII, H₂, and the number fraction of e^- (clockwise from upper left).

turned off, the gas quickly cools to $T \sim 10^3$ K through a combination of atomic, molecular hydrogen and Compton cooling. The molecular hydrogen formation time is shorter than the recombination time and a large amount of molecular hydrogen is produced, $x_{H_2} \sim \text{few} \times 10^{-3}$, irrespective of the density and temperature (see also Oh & Haiman 2002).² The pressure shock starts to dissipate and gas expelled from the center of the halo starts falling back in (for a more complete description, readers are encouraged to see MBH06).

Quantitatively, we concluded that the net impact of a transient UVB on star-formation could be understood simply in terms of the molecular hydrogen cooling time-scale:

$$t_{H_2} = \frac{1.5k_B T}{\Lambda_{H_2}} \frac{n_g}{n_{HI}n_{H_2}} \quad (1)$$

$$\approx \frac{1.5k_B T}{\Lambda_{H_2}} \frac{1}{n_g x_{H_2}} \quad (2)$$

$$\approx 4 \left(\frac{T}{10^3 \text{ K}} \right)^{-2.5} \left(\frac{x_{H_2}}{3 \times 10^{-3}} \right)^{-1} \left(\frac{n_g}{1 \text{ cm}^{-3}} \right)^{-1} \text{ Myr},$$

Here, k_B is the Boltzmann constant, T is the temperature, Λ_{H_2} is the H_2 cooling function, x_e is the free electron number fraction, n_g , n_{HI} , and n_{H_2} are the number densities of all baryons and electrons, neutral hydrogen, and H_2 , respectively, and the second equality is accurate shortly after $z_{UVB, \text{off}}$, since the recombination times at high redshifts and inside halos are very short.

² Glover & Abel 2008 have recently shown that including cooling due to excitations of H_2 through collisions with protons and electrons can lead to a lower molecular hydrogen fraction by a factor of 2. This would result in a longer delay in star-formation than shown here.

We found that the net *delay* in star-formation caused by the transient UVB is simply the ratio of the H_2 cooling time in the run with the UVB, to the H_2 cooling time in the run without UVB heating. This ratio is calculated near the halo core, shortly after the UVB disappears (when the shock first starts to dissipate and the gas behind it starts infalling again). For example, the delay in star-formation of a halo in the Heat0.8 run with respect to the same halo in the NoUVB run is:

$$f_{\text{delay}} \sim \frac{t_{H_2}^{\text{Heat0.8}}}{t_{H_2}^{\text{NoUVB}}} \sim \left(\frac{T^{\text{NoUVB}}}{T^{\text{Heat0.8}}} \right)^{2.5} \left(\frac{n_g^{\text{NoUVB}}}{n_g^{\text{Heat0.8}}} \right) \left(\frac{x_{H_2}^{\text{Heat0.8}}}{x_{H_2}^{\text{NoUVB}}} \right)^{-1} \quad (3)$$

We will now demonstrate that eq. (3) is equally adept at quantifying feedback in our new simulations. First, we show the spherically averaged radial profiles of the same (typical) halo in the NoUVB (solid lines), Heat0.08 (dashed lines) and Heat0.8 (dotted lines) simulation runs in our Figure 1. This halo, which has a mass of $4 \times 10^5 M_\odot$ at $z = 24.62$, was first able to form CD gas at $z = 20, 21, 19$ in the NoUVB, Heat0.08, and Heat0.8 runs, respectively. The top and bottom pairs of figures are snapshots at $z = 24.62, 23, 21, 19$ (left to right). The *upper panels* show the hydrogen density, mass-weighted gas temperature, gas cooling time, and radial velocity (clockwise from upper left). The *bottom panels* show mass fractions of HI, HII, H₂, and the number fraction of e^- (clockwise from upper left).

The profiles confirm the qualitative story line from the beginning of this section. The outward-moving pressure shock is evident in both UVB runs from the density and velocity panels at $z = z_{UVB, \text{off}}$, just before the UVB was turned off. This gas has already cooled to $T = 10^3$ K and started to collapse back onto the halo only a few Myr afterward at $z = 23$. Note that the outflow

and the resulting suppression of gas near the core is stronger in the run with a stronger UVB. Halos in the runs with a UVB also experience a strong boost in the H_2 fraction following $z_{UVB,off}$, with the increase being insensitive to the strength of the UVB. Formation of CD gas in this halo is notably delayed until $z = 19$ in the Heat0.8 run, while being mildly expedited in the Heat0.08 run. Can this behavior be understood through eq. (3)?

Let us focus on the output we have shortly after $z_{UVB,off}$, at $z = 23$. We can note that at this redshift, the pressure-driven shock has almost dissipated and reached zero velocity with gas recommencing infall behind the shock. Plugging in appropriate values into eq. 3 from the inner annulus of the halo, we obtain values of $f_{delay} \approx 0.5$ and 1.5 for the Heat0.08 and Heat0.8 runs, in good agreement with the simulation outcomes. This behavior again supports our conclusions in MBH06, where the same analysis held for a different, randomly-chosen halo, and implies that we understand the physical picture associated with feedback in relic HII regions.

In order to quantify the suppression of CD gas by a transient UVB on our entire sample of halos, we define (as in MBH06) the cumulative, fractional suppression of the number of newly formed halos as

$$\delta_{N,cd}(z) \equiv \frac{N_{cd}^{runi}(z) - N_{cd}^{runi}(z_{UVB,on})}{N_{cd}^{NoUVB}(z) - N_{cd}^{NoUVB}(z_{UVB,on})} - 1, \quad (4)$$

where $N_{cd}^{NoUVB}(z)$ and $N_{cd}^{runi}(z)$ are the total number of halos with CD gas at redshift z in the NoUVB run and some given run i , respectively. This expression is well-defined for $N_{cd}^{NoUVB}(z) > N_{cd}^{NoUVB}(z_{UVB,on})$; for $N_{cd}^{NoUVB}(z) = N_{cd}^{NoUVB}(z_{UVB,on})$, we set $\delta_{N,cd}(z) \equiv 0$. Note that by definition, $N_{cd}^{NoUVB}(z_{UVB,on}) = N_{cd}^{runi}(z_{UVB,on})$.

Similarly, we define the cumulative, fractional suppression of the CD gas mass as

$$\delta_{M,cd}(z) \equiv \frac{M_{cd}^{runi}(z) - M_{cd}^{runi}(z_{UVB,on})}{M_{cd}^{NoUVB}(z) - M_{cd}^{NoUVB}(z_{UVB,on})} - 1, \quad (5)$$

where $M_{cd}^{NoUVB}(z)$ and $M_{cd}^{runi}(z)$ are the total mass of CD gas at redshift z in the NoUVB run and some given run i , respectively. The total CD gas mass is obtained by merely summing the CD gas masses for all of the halos in the simulation. As for equation (4), this expression is well-defined for $M_{cd}^{NoUVB}(z) > M_{cd}^{NoUVB}(z_{UVB,on})$, and for $M_{cd}^{NoUVB}(z) = M_{cd}^{NoUVB}(z_{UVB,on})$, we set $\delta_{M,cd}(z) \equiv 0$.

Equations (4) and (5) provide an estimate of how the CD gas has been affected by the presence of a UVB, *following* the turn-on redshift of the UVB, $z_{UVB,on}$ (the values at $z_{UVB,on}$ are subtracted in order to provide a more sensitive measure of *relative* changes of CD gas). As defined above, $\delta_{N,cd}(z) = 0$ and $\delta_{M,cd}(z) = 0$ if the UVB has no effect. If the effect of a UVB is positive, resulting in positive feedback, $\delta_{N,cd}(z)$ and $\delta_{M,cd}(z)$ would be positive. If the effect of the UVB is negative, $\delta_{N,cd}(z)$ and $\delta_{M,cd}(z)$ would be negative.

In Figure 2, we plot the values of $M_{cd}^{runi}(z)$ (*top panel*) and $N_{cd}^{runi}(z)$ (*bottom panel*) in our four simulation runs with no LW background: runi = NoUVB (crosses), Flash (dashes), Heat0.08 (triangles), and Heat0.8 (squares). We note that the number of halos with CD gas increases in all runs until $z \sim 15$, when our central refined regions starts becoming more non-linear. At these lower redshifts, halos become contaminated with large DM particles entering from outside the refined region and we remove them from our analysis sample (see §2).

The corresponding values of $\delta_{M,cd}(z)$ and $\delta_{N,cd}(z)$ are plotted in Figure 3 in the top and bottom panels, respectively. Although




Figure 2. Values of $M_{cd}^{runi}(z)$ (*top panel*) and $N_{cd}^{runi}(z)$ (*bottom panel*) as defined in equations (5) and (4). The results are displayed for the NoUVB (crosses), Flash (dashes), Heat0.08 (triangles), and Heat0.8 (squares) simulation runs.

some of the notable fractional changes shown in Figure 3 might appear statistically insignificant due to the small number statistics inferred from Figure 2, it should be noted that these runs are not uncorrelated experiments. In other words, each of our runs in Table 1 is seeded with the same initial conditions, and so small relative changes compared to the NoUVB run are significant (i.e. the errors are not Poisson). Nevertheless, there are some minor differences between the runs in the amount of “contamination” by DM particles outside of our refined region.

In MBH06, we presented two main inferences regarding the effects of a transient UVB (aside from the delay due to cooling times discussed above): (1) a “critical” background value of $J_{21}^{UV} \sim 0.1$ separated positive from negative feedback (stronger backgrounds result in negative feedback, while weaker backgrounds result in positive feedback; see the discussion surrounding eq. 3 above); (2) the feedback effects, regardless of the sign, were transient and started to “disappear” by $z \sim 18$. However, we were unable to evolve our simulations below $z < 18$, since the central refined region became too non-linear and further progress was computationally prohibitive. Thus point (2) was an inference based on a figure analogous to Fig. 3.

In this work we present results from a less biased region and are able to go to lower redshifts, $z \sim 13.5$. Thus we can test both of the conclusions above and see if they are sensitive to the large-scale halo environment and redshift. From figure 3, we see that *both* conclusions are valid for these simulations as well. In other words, our transient (corresponding to the expected lifetime of a Pop III star) UVB results in transient feedback which disappears after $\sim 30\%$ of the Hubble time, with $J_{21}^{UV} \sim 0.1$ separating net positive and negative feedback. *Thus, these conclusions seem to be fairly insensitive to the large-scale overdensity of the simulated region, and by extension to the redshift-dependent properties of halos at $z_{UVB,on}$.*




Figure 3. Values of $\delta_{M,cd}(z)$ (top panel) and $\delta_{N,cd}(z)$ (bottom panel) as defined in equations (5) and (4). The results are derived from Figure 2 and displayed for the Flash (dashes), Heat0.08 (triangles), and Heat0.8 (squares) simulation runs.

It is important to note a new result evident in Fig. 3: *all runs with a transient UVB experience eventual positive feedback at $z \lesssim 15$* . This is true regardless of the strength of the UVB. We will examine this interesting result further in §4.

3.1 Adding a Persistent LW Background

Next, we investigate how the combination of transient UV heating and a persistent LW background impact star formation. This is a more physically-relevant scenario, since a LW background is likely established before a large fraction of the Universe has been reionized (Haiman et al. 2000).

Photons in the LW band dissociate H_2 molecules; thus by definition they provide additional negative feedback, undercutting some of the enhancement in the H_2 abundances in relic HII regions. Negative feedback from a LW background kicks in when the H_2 dissociation timescale becomes shorter than the H_2 formation timescale. Since the formation timescale is inversely proportional to gas density, whereas the dissociation timescale is independent of density, the density decrease caused by UV heating should make halos more susceptible to the negative feedback of a LW background (Oh & Haiman 2003)³. We investigate these processes further below.

Specifically, in Fig. 4, we plot values of $\delta_{M,cd}(z)$ (top panel) and $\delta_{N,cd}(z)$ (bottom panel) for our runs including a LW background. The LW background turns on at $z = 24.62$ and remains on. At the simplest level, we can see the suppression of molecular hydrogen (and hence cooling) due to the LW background by

³ Note that the temperature increase associated with a UVB does not contribute much to this combined negative feedback because the ionized gas is able to Compton cool down to the virial temperatures of these minihalos.




Figure 4. Suppression of cold dense gas in halos in simulations runs that include a persistent LW background. The LW background had specific intensities of $J_{LW} = 10^{-4}, 10^{-3}, 0.01$, or 0.1 (normalized at 12.87 eV, in units of $10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$). Each of these three LW backgrounds is applied to both our NoUVB and Heat0.8 runs at $z = 24.62$, and is subsequently left on. Values of $\delta_{M,cd}(z)$ (top panel) and $\delta_{N,cd}(z)$ (bottom panel) are shown, as defined in equations (5) and (4).

the more negative values of both $\delta_{M,cd}(z)$ and $\delta_{N,cd}(z)$ at fixed redshift.

Does this panel show evidence for the *additional* negative impact of the transient UVB discussed above? Without UV heating, a LW background with a specific intensity of $J_{21}^{LW} = 0.01$ is needed for notable negative feedback (see the blue empty triangles at $z \leq 14$ in the lower panel). This value is similar to the one found in MBH06, where we showed that by equating H_2 -cooling and H_2 -photodissociation one expects this critical LW background to scale as $J_{21}^{LW} \propto n_g f_{H2}/T$.⁴ When the UV heating is added, this critical value of J_{21}^{LW} causing negative feedback decreases by a factor of ~ 10 to $J_{21}^{LW} \sim 10^{-3}$ (see the green solid circles in lower panel at $z \leq 14$). This decrease seems to confirm the above arguments. However, such an interpretation is too simplistic and the negative impact of the LWB can be ameliorated by positive feedback inside the HII region at lower redshifts, as we shall see in §4.

From Fig. 4 one can also note that a value of $J_{21}^{LW} \sim 10^{-3} - 10^{-2}$ separates feedback regimes dominated by a LW background from those dominated by our transient UVB. This can be seen by the fact that the amount of suppression differs between the NoUVB and Heat0.8 cases at low values of the LW background (i.e. for $J_{21}^{LW} < 10^{-3}$), while at large values of J_{21}^{LW} , the amount of suppression is independent of the UVB. Near the threshold value of J_{21}^{LW} , negative feedback transitions from being UV heating dominated (at $z \gtrsim 17$, or $\lesssim 100$ Myr after $z_{UVB,off}$) to being LW

⁴ Since the mean density of gas collapsing into halos in these simulations is less than a factor of two smaller than those in MBH06, we roughly expect this critical value of J_{21}^{LW} to also be less than a factor of two smaller (at least in the linear regime).

background dominated ($z \lesssim 17$, or $\gtrsim 100$ Myr after $z_{\text{UVB,off}}$). As highlighted in MBH06, this “transition” behavior can be understood as the combined result of two effects: the UV heating is turned off, and its impact is transient, while the critical LW background scales roughly with the inverse of the density (Haiman et al. 2000; Oh & Haiman 2003) and hence a fixed LW background will have a larger impact at lower densities or decreasing redshifts.

4 EVENTUAL POSITIVE FEEDBACK

So far, we have not focused on the interesting new result hinted by the previous figures (e.g. see the $z \lesssim 14.5$ points in Fig. 3). This figure shows evidence of delayed (approaching our lowest redshifts) *positive* feedback in *all* runs with a UVB. This result is somewhat surprising given that the Heat0.8 run shows strong negative feedback initially after $z_{\text{UVB,off}}$. In fact, as was already pointed out when discussing Fig. 3, the initial feedback, whether positive or negative, fades away in *all* runs by $z \sim 17$.

In Fig. 5, we show the total gas fractions (*top panels*) and CD gas fractions (*bottom panels*) from halos at $z = 13.5$. The positive feedback mentioned above is immediately evident from this figure, and shows explicitly the masses of the affected halos. In particular, the NoUVB run at $z = 13.5$ has 3 *fewer* halos hosting CD gas than the Flash and Heat0.08 runs, and 2 *fewer* than even the Heat0.8 run (see also Fig. 2). These are some of the smallest and youngest of such halos at these redshifts. We also verify that these halos formed in less biased and less overdense regions than the halos which didn’t experience such positive feedback.

What is the physical cause of such delayed positive feedback? Before we attempt to answer this question, let us briefly review the sequence of feedback-related events arising from a transient UVB:

- (i) When the UVB turns on, gas gets ionized and heated to temperatures $\gtrsim 10^4$ K.
- (ii) The temperature increase sets-off an outward moving pressure shock in the cores of halos, where density profiles have already steepened as gas approached hydrostatic equilibrium. This pressure shock smooths out the gas distribution and leads to a decrease in gas density in the cores of halos.
- (iii) Once the UVB is turned off, the gas rapidly cools to $T \sim 10^3$ K through a combination of atomic, molecular hydrogen and Compton cooling. This temperature approximately corresponds to the gas temperature at the virial radius of such a proto-galactic, molecularly-cooled halo, thus effectively neutralizing the impact of temperature change on feedback.
- (iv) A large amount of molecular hydrogen is produced, $x_{\text{H}_2} \sim \text{few} \times 10^{-3}$, irrespective of the gas density and temperature.
- (v) The pressure-shock begins to dissipate and gas with a newly enhanced H_2 abundance starts falling back onto the partially evacuated halo.

It is the balance between the density depletion in step (ii) (which is dependent on the UVB intensity) and the enhanced cooling capability of the gas from step (iv) (which is independent of the UVB intensity), which determines the transient feedback outcomes discussed in § 3. But what happens to gas which is not in the cores of already-formed halos with a steepened density profile? In the extreme case of homogeneous gas, the UVB-induced temperature increase in (i) *does not translate into a pressure shock*. Thus, there is no density depletion from step (ii). In the more realistic case of a clumpy gas distribution, pressure gradients will still

form and smooth out the gas distribution, but the resulting depletion will be less than in the cores of halos in hydrostatic equilibrium. However, such lower-density gas outside of halos will still be able to cool to temperatures of $\sim 10^3$ K since the Compton cooling time is independent of density, and at these high redshifts is over a factor of two shorter than the recombination time at mean density [in fact, these lower density regions take longer to recombine and hence can Compton-cool for a longer period reaching lower temperatures (Oh & Haiman 2002); see also equations (2) and (3) in MBH06]. Likewise, step (iv) mentioned above also applies to such lower density gas, with the transient background resulting in an enhanced molecular hydrogen abundance. Thus, we expect gas which has not yet formed a self-similar (e.g. Navarro et al. 1996) halo profile to experience weaker negative feedback [i.e. step (ii)], and the same amount of positive feedback [i.e. step (iv)], when exposed to identical transient UVBs.

This conclusion is perhaps a bit counterintuitive, since many cosmological radiative feedback studies find that feedback is a strong function of halo mass, with more massive, earlier forming halos being *less* susceptible to negative feedback (Efsthathiou 1992; Barkana & Loeb 1999; Gnedin 2000; Shapiro et al. 2004; Kitayama & Ikeuchi 2000; Dijkstra et al. 2004; Mesinger & Dijkstra 2008). However, there are two important distinctions here: we are discussing feedback resulting from a *transient* UVB; and we are comparing halos which have not yet formed and set up a self-similar gas profile at $z_{\text{UVB,off}}$ to those which have already formed, rather than comparing *within* each of these two groups.

We also see some evidence of these trends in Fig. 6, where we plot the ratio (in runs with a UVB over the fiducial NoUVB run) of the H_2 fractions (*red squares*), the temperatures (*blue triangles*), and the average gas densities (*black circles*) within the central 15pc (proper) of halos. These quantities are shown shortly after $z_{\text{UVB,off}}$, at $z = 23$. As already noted in §3 and in (iv) above, the H_2 fraction tends to increase by the same factor, regardless of the nature of the heating. As a result, the sign of the overall feedback (negative or positive) is determined primarily by whether or not the depression of the gas density overcomes the positive effects of this H_2 enhancement. The density depletion does depend strongly on the type and amount of heating. This depletion, shown in black circles, is of course stronger for the Heat0.8 than the Heat0.08 run. Thus halos in the former initially experience strong negative feedback and in the latter negligible or weak positive feedback, as we’ve seen in §3 and as noted by the green crosses in the figure. Note however that the difference in the amount of gas depletion between these two heating runs decreases with decreasing halo mass as one begins being affected by Jeans/filtering scale effects⁵. Following the above argument, this trend should continue and density depletion should decrease in regions which are less evolved (i.e. the black points should approach unity as the halo mass is decreased below the values shown in the figure). It is these regions which become the progenitors of our halos experiencing eventual positive feedback. Unfortunately, at $z = 23$, these regions are too unevolved to be identified by our halo finding algorithm and thus do not appear in Fig. 6.

This eventual positive feedback should be quite sensitive to the LWB. Indeed, looking at the bottom panel of Fig. 4, we see

⁵ Note that accurately modeling Jeans smoothing requires very high resolution in both dark matter and gas (Naoz et al. 2009). These small scales are shown in Fig. 6 merely to show qualitative trends.




Figure 5. Total gas fractions (*top panels*) and cold, dense gas fractions (*bottom panels*) as a function of total halo mass at redshift $z = 13.5$, the lowest redshift output of our simulations. The four panels correspond to the NoUVB (*top left*), Flash (*top right*), Heat0.08 (*bottom left*), and Heat0.8 (*bottom right*) simulation runs. Note that all runs with a UVB have more halos with CD gas than the NoUVB run.

that this feedback effect is wiped-out if $J_{21}^{\text{LW}} \gtrsim 10^{-3}$. It is likely that such modest values of J_{21}^{LW} are reached well before the majority of the Universe is ionized (Haiman et al. 2000), though the buildup of the LWB can be slowed by an increasing optical depth to LW photons from the enhanced H_2 abundances in relic HII regions (Johnson et al. 2007), or in shells surrounding HII regions (Ricotti et al. 2001; Kuhlen & Madau 2005; though see Kitayama et al. 2004). Thus studying the true physical importance of this positive feedback mechanism would have to be done with self-consistent cosmological simulations of reionization.

Now, let us look at specific examples by studying the evolution of one such halo experiencing eventual positive feedback. In Figure 7, we plot the same quantities as in Fig. 1, but for a halo

which experiences positive feedback in the Heat0.8 run. Note that, unlike the more evolved halos which are subjected to the transient feedback and have already formed at $z_{\text{UVB,off}}$, the earliest snapshot we have of this halo is at $z = 21$. Solid and dashed curves correspond to the NoUVB and Heat0.8 runs, respectively.

It is important to note that this halo forms considerably after $z_{\text{UVB,off}}$, and thus the “density depletion” seen in the figure at $z \gtrsim 17$ is of a different nature than discussed in §3 and Fig. 1. Namely, this gas wasn’t evacuated from the halo center with a pressure shock. Instead, these late forming halos start pulling in baryons when their virial temperature exceeds that of the gas temperature. The heating and subsequent Compton cooling following the transient UVB sets the IGM gas temperature in these runs at $T \sim 10^3$ K. Thus baryonic infall onto newly formed halos can commence




Figure 6. Ratio of the H_2 fractions (red squares), the temperatures (blue triangles), and the average gas densities (black circles) within the central 15pc (proper) of halos. Ratios of these quantities in the Heat0.08 and Heat0.8 runs over the fiducial NoUVB run are shown in the left and right panels, respectively, at $z = 23$. The delay factors from eq. (3) are also plotted as green crosses.

when $T_{\text{vir}} \gtrsim 10^3$ K. Since the IGM is cooler in the NoUVB run, gas there gets a “head-start” in accreting onto DM halos. However, as is evident in the figure, the enhanced H_2 cooling channel in the Heat0.8 run allows the gas to quickly catch-up and surpass its twin in the NoUVB run. Thus by redshift $z = 14$, the halo has formed CD gas in the Heat0.8 run, but not in the NoUVB run.

How does adding a LWB impact this positive feedback? From Fig. 4, we have already surmised that a background intensity of $J_{21}^{\text{LW}} \gtrsim 10^{-3}$, can offset these trends by decreasing the enhanced H_2 fraction in relic HII regions. In Fig. 8, we plot profiles from the same halo in the $\text{LW}10^{-3}$ _NoUVB (solid curves) and $\text{LW}10^{-3}$ _Heat0.8 (dashed curves). By comparing the solid curves from Fig. 7 to the dashed curves from Fig. 8, one confirms that a LWB with $J_{21}^{\text{LW}} \sim 10^{-3}$ effectively neutralizes the positive feedback from the transient UVB. Thus at $z = 14$ the halo core has a similar gas profile and cooling time in the NoUVB and $\text{LW}10^{-3}$ _Heat0.8 runs.

By comparing the two profiles at $z = 14$ in Fig. 8, we see that the transient UVB still stimulates positive feedback, but the effect is not nearly as dramatic. Specifically, in the halo core at $z = 14$ the density is enhanced and the cooling time is lowered (both by a factor of ~ 2) in the $\text{LW}10^{-3}$ _Heat0.8 run compared to the $\text{LW}10^{-3}$ _NoUVB run. This level of positive feedback is much more modest compared with the case with no LWB, where the equivalent density enhancement was a factor of ~ 50 and the cooling time was lower by a factor of ~ 15 (see Fig. 7).

Does this trend continue as we increase the strength of the UVB? In Fig. 9, the solid and dashed curves correspond to the $\text{LW}10^{-2}$ _NoUVB and $\text{LW}10^{-2}$ _Heat0.8 runs at $z = 14$, respectively; similarly in Fig. 10, the solid and dashed curves correspond to the $\text{LW}0.1$ _NoUVB and $\text{LW}0.1$ _Heat0.8 runs at $z = 14$, respectively. Here we see that the core density in the $\text{LW}0.1$ _Heat0.8 run is lower than in the $\text{LW}0.1$ _NoUVB run, but the cooling time is also lower by the same factor of ~ 2 seen in Fig. 8, sourced by

the enhanced H_2 fraction. In fact, from higher redshift output and the previous figures, we see that the density is still “catching up” as time progresses; namely, the dashed density curve keeps getting closer to the solid density curve. Thus we see that *halos inside relic HII regions, when exposed to the same LWB, always have a higher H_2 abundance and shorter cooling times than halos outside relic HII regions, allowing gas to cool faster once it finally begins to collapse onto the halo.*

It is important to note that in this regime where a strong LWB dominates the overall feedback, star formation is likely to be substantially delayed. This delay can be long enough to allow the halo to be photoevaporated during reionization, before it had a chance to host additional stars. Thus if one is only concerned with the fate of those minihalos contributing their ionizing photons to the progress of reionization, it is possible that the sole *persistent* feedback mechanism is *positive*.

5 THE IMPACT OF NEGLECTING RADIATIVE TRANSFER

As we have noted a number of times in this paper, our simulations assume the optically thin limit, for both ionizing radiation and photodissociating LW photons. Both of these approximations clearly fail at points in our simulations, and so it is necessary to understand the nature and implications of this approximation. For clarity, we discuss ionizing photons and LW photons separately.

We discuss the impact of radiation hydrodynamic (RHD) effects due to ionizing photons first. There is now a substantial body of literature on the ionization of primordial gas halos (Shapiro et al. 2004; Whalen et al. 2004; Iliev et al. 2005; Alvarez et al. 2006; Susa & Umemura 2006; Whalen et al. 2008; Abel et al. 2007; Wise & Abel 2008b) and the broad outline of how halos are ionized is now well understood. We begin by noting that the optical depth of the halos which are ionized in our simulation




Figure 7. The same quantities as in Fig. 1, but for a halo which experiences positive feedback in the Heat0.8 run. Solid and dashed curves correspond to the NoUVB and Heat0.8 runs, respectively.




Figure 8. Same as Fig. 7, but with solid and dashed curves corresponding to the LW10⁻³_NoUVB and LW10⁻³_Heat0.8 runs, respectively.

box at $z = 25$ (as shown in Fig. 1), as measured from the virial radius to the center is – at the Lyman limit – well over 100. Therefore, at first blush, the optically thin limit seems like a poor approximation. However, this impression is not entirely accurate for two reasons.

First, we are studying the *later* formation of halos in relic HII regions, and much of the gas that eventually falls into a halo at late times is *not* in the center of halos at reionization. This is the question that Oh & Haiman (2003) asked: does the excess entropy generated in low-density gas in relic HII regions prevent it from forming stars at later times. This low density gas has much lower

optical depth and so is very likely to be ionized by a nearby Pop III star.

Second, even for the gas in the centers of our halos at $z_{\text{UVB, on}}$, the optically thin approximation is not as bad as the large optical depth at the Lyman limit indicates. We can see this more clearly by comparing our results to RHD calculations. For example, take the (typical) halo shown in Fig. 1. The central density when ionization begins (at $z = 25$) is just slightly over 1 cm^{-3} . This value is much below the central density (10^4 cm^{-3}) assumed by Susa & Umemura (2004), and is not that different from their critical density for self shielding (depending on the assumed distance




Figure 9. Same as Fig. 7, at $z = 14$, but with solid and dashed curves corresponding to the LW10 $^{-2}$ _NoUVB and LW10 $^{-2}$ _Heat0.8 runs, respectively.

of the source). A better comparison is with the “023” series of simulations run in Whalen et al. (2008), which have almost identical central densities. As is shown in Fig. 17 of that paper, this halo is disrupted and dissociated for all assumed distances of the first star. Comparing Fig. 9 and 10 of that paper to our Fig. 1 shows some obvious differences due to radiative transfer effects, but the basic evolution is not all that different. In particular, after a few Myr, the density at the halo center has decreased by about a factor of ~ 5 after ~ 3 Myr, while the molecular hydrogen fraction has increased to a few times 10^{-3} . These numbers are very similar to what we find, and it is these basic facts which, we argue, cause the delay in future star formation, as discussed earlier. The reason for the similarity is that, as it sweeps over the halo, the ionization front can quickly ionize hydrogen until it transitions to a D-type front and stalls. However, this happens only ~ 15 pc from the core, at den-




Figure 10. Same as Fig. 7, at $z = 14$, but with solid and dashed curves corresponding to the LW0.1_NoUVB and LW0.1_Heat0.8 runs, respectively.

ties slightly below 1 cm^{-3} , and the shock generated by the D-type front quickly disperses the core.

Of course, we do not claim to reproduce the radiation hydrodynamics in detail. For example, we do not reproduce the thin self-shielded filament at the lower right of Fig. 10 of Whalen et al. (2008), nor do we follow the detailed shock-compression or “rocket” offset of the core. However, by choosing to apply the ionizing flux at a redshift when the halos have low masses and central densities, we minimize the impact of the radiation transfer. To verify that this is actually true in our simulations, we examined the central densities and masses of our halos at $z = 25$; we find that all halos except for one have masses below $10^6 M_{\odot}$ and central densities below $\sim 2 \text{ cm}^{-3}$ (one halo has $M > 10^6 M_{\odot}$ and $n_c = 7 \times 10^3 \text{ cm}^{-3}$). Our halo abundance is consistent with predictions from the conditional Press-Schechter formalism (MBH06). We chose this approach because we are interested in investigating the impact of relic HII regions on the formation of structure which

forms significantly later, rather than studying the impact of a Pop III star on a nearby, highly collapsed halo. In particular, as noted in the introduction, our motivation is to follow up on the suggestion of Oh & Haiman (2003) that relic HII regions will imprint an excess entropy on the low-density gas at early times, which will then suppress later star formation. Of course, we stress that it is still important to examine our results with a more realistic RHD treatment.

The results discussed in the section on positive feedback (section 4) are even more robust because in that case the halos form out of gas that was barely overdense when the ionization occurred. As can be seen in Fig. 7, even at $z = 21$ (long after the transient ionization), the central density of the halo (in the noUVB case) is only 0.2 cm^{-3} .

Another useful comparison of our results can be made with Wise & Abel (2008a), who used a full cosmological simulation of Pop III star formation including radiative transfer (albeit in a region considerably smaller than considered here). While a direct comparison is difficult because Wise & Abel (2008a) do not explicitly measure the suppression due to photo-ionization, a number of conclusions can be drawn. First, it is clear from Fig. 7 of that paper that they also see the evaporation of gas from low-mass halos and filaments, and the resulting density field in the two simulations looks remarkably similar. In addition, the time difference between the first Pop III object and the formation of the second is about 80 Myr, consistent with the time-delay derived in this paper. On the other hand, one important point that comes out of that paper is that, because Pop III star formation tends to be clustered, relic HII regions do not sit isolated for longer periods of time and so a region of space may undergo multiple ionization episodes, something which is not considered in this work.

Next, we turn to LW photons. First, we remind the reader that we are discussing a persistent LW background, and that although the LW flux is produced by the same *class* of objects that generate the transient UV flux, it is not produced by the same physical objects. As discussed in more detail in Haiman et al. (1996), the LW background is accumulated from a large number of distant sources. The effective optical depth in the LW lines in the background IGM, over a Hubble distance, is at most $\tau \approx 1 - 2$ (Haiman et al. 2000; Ricotti et al. 2001), so the Olbers' integral for the LWB is dominated by the large number of sources at a good fraction of the Hubble distance. In comparison, the ionizing photons come exclusively from the much fewer source(s) nearby, within the local HII region.

LW photons can be self-shielded by a sufficient column of H_2 . Here again, there are two regimes. One is the large quantities of low-density gas in the relic HII regions that form substantial amounts of molecular hydrogen immediately after the ionizing flux is turned off (e.g. Kuhlen & Madau 2005). Although H_2 formed in this manner may slow the build-up of a persistent LW background, it does not – for a given LW background – provide a large amount of shielding. For example, Johnson et al. (2007) find typical values of LW optical depths of order unity for a single relic HII region. This means that the LW flux inside the relic HII region can be reduced by a factor of a few, but it also means that it doesn't take long for the persistent LW background to "burn off" the H_2 produced in relic HII regions.

The second regime occurs after the ionized gas falls back into the collapsed halos, where H_2 is in equilibrium between formation and photo-destruction. In this case, when a sufficient column density of H_2 is formed, it can effectively attenuate the LW flux. Based on our simulations, we find that when CD gas forms, the LW optical depth can be of order unity or larger, indicating that self-shielding can play a role during the recollapse (although probably not during

the photoionization process itself). Therefore, we caution readers that the critical LW values we give above may need to be increased when self-shielding is properly accounted for.

Finally, we note that in simulations with full radiative transfer, the photoionizing flux can also vary across a given halo (relic H II regions have radii of 2-3 kpc, while the Lagrangian radius of a typical halo in our simulations is of order 0.5 kpc). This also means that, while most forming halos lie either completely inside or outside a relic H II region, some proto-halos could be partially ionized.

6 CONCLUSIONS

UV radiation from early astrophysical sources could have a large impact on subsequent star formation in nearby protogalaxies, and in general on the progress of cosmological reionization. Theoretical arguments based on the absence of metals in the early Universe suggest that the first stars were likely massive, bright, yet short-lived, with lifetimes of a few million years. Here we study the impact of the transient radiation arising from such stars on early protogalaxies. We apply the same statistical approach as in MBH06, studying various combinations of transient UVBs and persistent LWBs, using the hydrodynamical simulation code, Enzo. We also study a more typical and relevant region whose proto-galaxies form at lower redshifts, $z \sim 13-20$. This allows us to trace feedback effects on longer time-scales and lower redshifts.

We confirm the results of MBH06 that feedback in the relic HII regions resulting from such transient radiation, is itself transient. Feedback effects dwindle away after $\sim 30\%$ of the Hubble time, and the same critical specific intensity of $J_{\text{UV}} \sim 0.1 \times 10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ separates positive and negative feedback regimes. A weaker UVB stimulates subsequent star formation inside the relic HII regions, by enhancing the molecular hydrogen abundance. A stronger UVB delays star-formation by reducing the gas density at the centers of collapsing halos. The fact that we have confirmed the results in MBH06 (which were mostly inferences obtained by extrapolating from higher redshifts since we were unable to continue those simulations to low enough redshifts) suggests that overall feedback is fairly insensitive to the large-scale environment, overdensity, and redshift-dependent halo parameters, and can accurately be modeled in this regime with just the intensity of the impinging UVB.

We note that a value of $J_{21}^{\text{LW}} \sim 10^{-3}-10^{-2}$ separates feedback regimes dominated by the LW background from those dominated by the transient UVB. As discussed in section 5, there is some uncertainty in these values due to our optically thin treatment. In particular, accounting for LW self-shielding may increase the values of J_{21}^{LW} quoted here.

Additionally, we discover a second episode of eventual positive feedback in halos which have not yet collapsed when their progenitor regions were exposed to the transient UVB. Gas in such late-forming objects did not have time to set-up a steep density profile, which would translate into a strong pressure shock when photo-heated. Instead they formed out of material which was of more moderate densities at $z_{\text{UVB, on}}$ where the lasting impact of the transient UVB was positive: seeding the gas with an enhanced abundance of molecular hydrogen, which aids in its eventual cooling. However, this feedback regime is very sensitive to the presence of Lyman-Werner radiation, and disappears under fairly modest background intensities of $J_{\text{LW}} \gtrsim 10^{-3} \times 10^{-21} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$. Nevertheless, when exposed to

the same LWB, halos inside relic HII regions always have a higher H_2 abundance and shorter cooling times than halos outside relic HII regions, allowing gas to cool faster once it finally begins to collapse onto the halo. Although it is difficult to accurately estimate the build-up of the LWB, it seems likely that such modest values were surpassed well before the bulk of reionization (Haiman et al. 2000). Thus this persistent eventual positive feedback might only be of academic interest, provided LW self-shielding is not efficient.

We conclude that radiative feedback from a short-lived UVB seems unlikely to have a major impact on the progress of global cosmological reionization, provided that present estimates of the luminosities and lifetimes of Pop III stars are accurate (e.g. Schaerer 2002). Thus it is likely that the LWB plays the dominant role in regulating feedback in relic HII regions.

We thank the referee for comments which significantly improved the presentation of this paper. Support for this work was provided by NASA through Hubble Fellowship grant #HF-01222.01 to AM, awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555. GB acknowledges support from NSF grants AST-05-07161, AST-05-47823, and AST-06-06959, as well as computational resources from the National Center for Supercomputing Applications. ZH thanks the Polányi Program of the Hungarian National Office of Technology.

REFERENCES

- Abel T., Bryan G. L., Norman M. L., 2002, *Science*, 295, 93
 Abel T., Wise J. H., Bryan G. L., 2007, *ApJ*, 659, L87
 Ahn K., Shapiro P. R., 2007, *MNRAS*, 375, 881
 Alvarez M. A., Bromm V., Shapiro P. R., 2006, *ApJ*, 639, 621
 Barkana R., Loeb A., 1999, *ApJ*, 523, 54
 Bromm V., Coppi P. S., Larson R. B., 2002, *ApJ*, 564, 23
 Bryan G. L., 1999, *Comput. Sci. Eng.*, 46, 1
 Cen R., 2003a, *ApJ*, 591, L5
 Cen R., 2003b, *ApJ*, 591, 12
 Ciardi B., Ferrara A., Abel T., 2000, *ApJ*, 533, 594
 Dijkstra M., Haiman Z., Rees M. J., Weinberg D. H., 2004, *ApJ*, 601, 666
 Dunkley J., et al., 2008, *ArXiv e-prints*, 803
 Efsthathiou G., 1992, *MNRAS*, 256, 43P
 Eisenstein D. J., Hu W., 1999, *ApJ*, 511, 5
 Glover S. C. O., Abel T., 2008, *MNRAS*, 388, 1627
 Gnedin N. Y., 2000, *ApJ*, 542, 535
 Haiman Z., Abel T., Rees M. J., 2000, *ApJ*, 534, 11
 Haiman Z., Bryan G. L., 2006, *ApJ*, 650, 7
 Haiman Z., Holder G. P., 2003, *ApJ*, 595, 1
 Haiman Z., Rees M. J., Loeb A., 1996, *ApJ*, 467, 522
 Haiman Z., Rees M. J., Loeb A., 1997, *ApJ*, 484, 985
 Iliev I. T., Shapiro P. R., Raga A. C., 2005, *MNRAS*, 361, 405
 Johnson J. L., Greif T. H., Bromm V., 2007, *ApJ*, 665, 85
 Kitayama T., Ikeuchi S., 2000, *ApJ*, 529, 615
 Kitayama T., Yoshida N., Susa H., Umemura M., 2004, *ApJ*, 613, 631
 Komatsu E., et al., 2008, *ArXiv e-prints*, 803
 Kuhlen M., Madau P., 2005, *MNRAS*, 363, 1069
 Machacek M. E., Bryan G. L., Abel T., 2001, *ApJ*, 548, 509
 McGreer I. D., Bryan G. L., 2008, *ApJ*, 685, 8
 Mesinger A., Bryan G. L., Haiman Z., 2006, *ApJ*, 648, 835
 Mesinger A., Dijkstra M., 2008, *MNRAS*, 390, 1071
 Naoz S., Barkana R., Mesinger A., 2009, *MNRAS*, pp 1223–+
 Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563
 Norman M. L., Bryan G. L., 1999, in *ASSL Vol. 240: Numerical Astrophysics Cosmological Adaptive Mesh Refinement*. pp 19–+
 Oh S. P., Haiman Z., 2002, *ApJ*, 569, 558
 Oh S. P., Haiman Z., 2003, *MNRAS*, 346, 456
 O’Shea B. W., Abel T., Whalen D., Norman M. L., 2005, *ApJ*, 628, L5
 O’Shea B. W., Norman M. L., 2008, *ApJ*, 673, 14
 Ricotti M., Gnedin N. Y., Shull J. M., 2001, *ApJ*, 560, 580
 Ricotti M., Gnedin N. Y., Shull J. M., 2002, *ApJ*, 575, 49
 Schaerer D., 2002, *A&A*, 382, 28
 Schaerer D., 2003, *A&A*, 397, 527
 Shapiro P. R., Iliev I. T., Raga A. C., 2004, *MNRAS*, 348, 753
 Shapiro P. R., Kang H., 1987, *ApJ*, 318, 32
 Spergel D. N., Verde L., Peiris H. V., Komatsu E., Nolte M. R., Bennett C. L., Halpern M., Hinshaw G., Jarosik N., Kogut A., Limon M., Meyer S. S., Page L., Tucker G. S., Weiland J. L., Wollack E., Wright E. L., 2003, *ApJS*, 148, 175
 Susa H., Umemura M., 2004, *ApJ*, 600, 1
 Susa H., Umemura M., 2006, *ApJ*, 645, L93
 Whalen D., Abel T., Norman M. L., 2004, *ApJ*, 610, 14
 Whalen D., O’Shea B. W., Smidt J., Norman M. L., 2008, *ApJ*, 679, 925
 Wise J. H., Abel T., 2007, *ApJ*, 671, 1559
 Wise J. H., Abel T., 2008a, *ApJ*, 684, 1
 Wise J. H., Abel T., 2008b, *ApJ*, 685, 40
 Wyithe J. S. B., Loeb A., 2003, *ApJ*, 588, L69
 Yoshida N., Oh S. P., Kitayama T., Hernquist L., 2007, *ApJ*, 663, 687
 Yoshida N., Omukai K., Hernquist L., 2008, *Science*, 321, 669